



## A STUDY OF BRACHIAL ARTERY WITH HIGH UP DIVISION AND ITS CLINICAL SIGNIFICANCE

Jitendra Gupta\*, Ravi Jain, Manish Patil

Department of Anatomy, R. D. Gardi Medical College, Agar Road, Surasa, Ujjain-456006, India

\*Corresponding Author: Dr. Jitendra Gupta, Associate Professor, Department of Anatomy, R. D. Gardi Medical College, Ujjain, India

Received for publication: July 23, 2012; Accepted: September 14, 2012.

**Abstract:** Anomalies of the forelimb arterial tree are fairly common. In present study we observed the brachial artery with its high up division into radial and ulnar arteries in the middle of the arm. The findings were noted after thorough dissection of the upper limbs of both sides (axilla, arm, cubital fossa, forearm and palm) in twenty cadavers in the Department of Anatomy, R. D. Gardi Medical College, Ujjain, in the period of August 2010 to July 2012. After dissection each dissected part was numbered and arteries were painted and Photographic documentation of the variation was taken. An unusually short segment brachial artery was noted in the 2 cadaver. This short segment brachial artery bifurcated more proximally at the level of insertion of coracobrachialis in the middle of the left arm into radial and ulnar arteries. Further distribution of these two arteries was normal in both cadavers. No other variation was found particularly in relation to the cords of the brachial plexus and their branches. The short segment brachial artery with high up bifurcation as noted in the present study is fairly common. The variations can be explained in the light of embryological development. In addition, knowledge of such variation is important for carrying out surgical procedures in the arm in the arm and measurement of blood pressure routinely.

**Keywords:** Brachial Artery, Radial Artery, Ulnar Artery.

### INTRODUCTION

The brachial artery usually begins as a continuation of the axillary artery at the distal (inferior) border of the tendon of teres major and ends about a centimeter distal to the elbow joint (at the level of the neck of the radius) by dividing into radial and ulnar arteries [1].

Occasionally the artery divides proximally into two trunks, which may reunite. Frequently it divides more proximally than usual and this unusually short segment brachial artery may bifurcate as usual or it may trifurcate into radial, ulnar and common interosseous arteries [2]. Several other variations related to the termination of such a short segment brachial artery have been mentioned by some earlier workers.

Such variations can be explained on the basis of embryogenic development. According to Feinberg, ectodermal mesenchymal interactions and extracellular matrix components within the developing limb bud are controlling the initial patterning of blood vessels [3]. Further, there is a view that some inductive factors from the limb mesenchyme cause the changes in the blood vessel pattern [2]. High up division of the brachial artery can also be explained on the basis of observations made by Arey in 1957 where he highlighted that, there may be persistence of vessels which normally obliterate and disappearance or failure of development of vessels which normally persist [4].

This reversal of the normal process of vascular development is largely due to altered local hemodynamic environment [5]. Finally, knowledge of such variations has got clinical importance especially in the field of orthopedic, vascular and plastic surgeries [6].

The main objective of the present study is to document an unusually short segment brachial artery with its high up division into radial and ulnar arteries in the middle of the arm and to establish embryological basis and applied-anatomical correlations of such variation.

### MATERIALS AND METHODS


We conducted this study during routine dissection of the upper limbs of both sides (axilla, arm, cubital fossa, forearm and palm) in twenty cadavers in the Department of Anatomy, R. D. Gardi Medical College, Ujjain. After dissection each dissected part was numbered and arteries were painted and Photographic documentation of the variation was taken. The length of Brachial artery was measured in each specimen with the measuring tape from the lower border of teres major muscle to the bifurcation of brachial artery and reading was noted. (Table.1)



## RESULTS

Among twenty cadavers, two cadaver show high up division of brachial artery. In one cadaver of a 60year-old female, an unusually short segment brachial artery with high up division of brachial artery in the middle of the left arm was observed. The unusually short-segment brachial artery was 8.6cm in length and having slightly less caliber than usual. However it bifurcated normally into ulnar and radial arteries with same calibers. Further distribution of these two arteries was normal. Radial recurrent and common interosseous arteries arose normally from the radial and ulnar arteries, respectively. Profunda brachii and other branches of the brachial artery arose from it prior to its bifurcation. No other variation was found particularly in relation to the cords of the brachial plexus, their branches and their relations with axillary and brachial arteries. Dissection of the right upper limb revealed no unusual observations.

Another cadaver of 58years male was dissected and the high up division of brachial artery at the level of insertion of coracobrachialis in the middle of the Right arm was observed. The unusually short-segment brachial artery was 12.2cm in length and having same caliber as usual. However it bifurcated normally into ulnar and radial arteries with same calibers. Further distribution of these two arteries was normal. Radial recurrent and common interosseous arteries arose normally from the radial and ulnar arteries, respectively. Profunda brachii and other branches of the brachial artery arose from it prior to its bifurcation. No other variation was found particularly in relation to the cords of the brachial plexus, their branches and their relations with axillary and brachial arteries. Dissection of the left upper limb revealed no unusual observations. Finally an attempt was made to explain the abnormality in the light of embryological development and to establish its clinical implications.



**Table.1:** Length of Brachial artery in cadavers.

S.NO	Sex	Brachial artery length (cm)	
		Right	Left
1	Female	24.1	24
2	Male	23.8	23.8
3	Male	12.2	24.2
4	Female	22.1	22.3
5	Female	22.5	22.4
6	Male	21.9	22.2
7	Female	22.3	8.6
8	Male	21.4	21.3
9	Male	22.4	22.6
10	Male	23.4	23.8
11	Male	22.8	22.2
12	Male	24.0	24.2
13	Female	23.0	23.2
14	Female	22.8	22.4
15	Male	24.2	24.2
16	Male	21.8	22.2
17	Female	23.8	23.6
18	Male	21.8	22.0
19	Male	24.4	24.2
20	Female	22.0	21.6

## DISCUSSION

Anomalies of the forelimb arterial tree are fairly common. This is mainly because of their multiple and plexiform sources, the temporal succession of emergence of principal arteries, anatomises and periarticular networks and functional dominance followed by regression of some paths [2]. Occasionally the artery divides proximally into two trunks, which may reunite. Frequently it divides more proximally than usual, and this unusually short segment brachial artery may bifurcate as usual or it may trifurcate into radial, ulnar and common interosseous arteries. More often the radial branches arise proximally, leaving a common trunk for the ulnar and common interosseous; sometimes the ulnar artery arise proximally, the radial and common interosseous forming the other division; the common interosseous may also arise proximally [1].

Keen in 1961 mentioned high origin of radial artery from the brachial artery. He explained this variant on the basis of Arey's observations regarding anomalous blood vessels [4, 7]. In fact, Keen highlighted that there was persistence of the upper portion of the radial artery arising from the brachial artery proximal to the origin of ulnar artery followed by failure of development of the new connection of the radial artery with the brachial artery at the level of origin of ulnar artery [7]. Guha et al. observed high up division of brachial artery into radial and ulnar arteries in the middle of the arm associated with variant median nerve and absent musculocutaneous nerve [8].

The unusually short segment brachial artery with its high up division into radial and ulnar arteries as observed in the present study can be explained in the light of embryogenic development. The early limb bud receives blood via intersegmental arteries, which contribute to a primitive capillary plexus. At the tip of

the limb bud there is a terminal plexus that is constantly renewed in a distal direction as the limb grows. Later one main vessel supplies the limb and the terminal plexus; it is termed the axis artery. The aforesaid terminal plexus at the tip of the limb bud is separated from the outer ectodermal sleeve of the limb by an avascular zone of mesenchyme. This avascular region contains an extracellular matrix consisting largely of hyaluronic acid. Removal of this hyaluronic acid by hyaluronidase results in vascularization of the tissue since partial degradation products of hyaluronic acid are angiogenic. Thus ectodermal mesenchymal interactions and extracellular matrix components are controlling the initial patterning of blood vessels within the limb [3]. In the upper limb bud the axis artery is derived from the lateral branch of the seventh intersegmental artery (subclavian). The arterial trunk grows outwards along the ventral axial line and terminates in the deep plexus in the developing hand. Proximal part of the main trunk forms the axillary and brachial arteries and its distal part persists as the anterior interosseous artery and the deep palmar arch. The radial and ulnar arteries are the latest arteries to appear in the forearm from the axis artery (brachial). In the present case both the radial and ulnar arteries arose more proximally from the brachial artery leading to the formation of an unusually short segment and slightly narrow brachial artery which bifurcated into radial and ulnar arteries at a more proximal level near the middle of the arm. However, the subsequent distribution of the radial and ulnar arteries was normal. Campta highlighted diagnostic, interventional and surgical significance of such a variation. Diagnostically this type of variation may disturb the evaluation of angiographic images. Further knowledge of such variation has got clinical importance especially in the field of orthopedic, plastic and vascular surgeries [6].

### CONCLUSION

The short segment brachial artery and its variant termination in the form of high up bifurcation as noted in the present study are fairly common. The case can be explained in the light of embryological development. In addition, knowledge of such variation is important for carrying out surgical procedures in the arm and measurement of blood pressure routinely.

### ACKNOWLEDGEMENT

The authors are grateful to Dr. V. K. Mahadik, Director of R. D. Gardi Medical College, Ujjain for his advice and encouragement. Additional thanks go to Dr. Ashutosh Chourishi for editorials assistance.

### REFERENCES

1. Johnson D, Ellis H, Collins P, Upper Arm. In: Standing S, Ellis H, Healy JC, Johnson D, Williams A, Collins P, eds. Gray's Anatomy, 39th Ed., Philadelphia, Elsevier Churchill Livingstone, 2005, 856.
2. Williams PL, Bannister LH, Berry MM, Collins P, Dyson M, Dussek JE, Ferguson MW, eds. Gray's Anatomy. 38th Ed., London, Churchill Livingstone. 1999; 319, 1539.
3. Feinberg RN, Vascular development in the embryonic limb bud. In: Feinberg RN, Sherer GK, Auerbach R, eds. The development of the vascular system. Basel, Karger (Issues Biomed), 1991; 14: 136-148.
4. Arey LB, Developmental Anatomy, 6th Ed., Philadelphia, WB Saunders, 1957; 375-377.
5. Rodriguez-Baeza A, Nebot J, Ferreira B, Reina F, Perez J, Sanudo JR, Roig M, An anatomical study and ontogenetic explanation of 23 cases with variations in the main pattern of the human brachio-antebrachial arteries, J. Anat., 1995; 187: 473-479.
6. Gonzalez-Compta X, Origin of the radial artery from the axillary artery and associated hand vascular anomalies, J. Hand. Surg. Am., 1991; 16: 293-296.
7. Keen JA, A study of the arterial variations in the limbs, with special reference to symmetry of vascular patterns, Am. J. Anat., 1961; 108: 245-261.
8. Guha R, Palit S, A rare variation of anomalous median nerve with absent musculo cutaneous nerve and high up division of brachial artery, J. Interacad., 2005; 9: 398-403.

Source of support: Nil

Conflict of interest: None Declared