



A STUDY OF DYSFUNCTIONAL UTERINE BLEEDING: CLINICAL FACTORS AND ENDOMETRIAL HISTOLOGY

Seena KB

Department of Obstetrics & Gynecology, Government Medical College Hospital, Thrissur, Kerala, India

Received for publication: February 21, 2015; Revised: February 27, 2015; Accepted: March 10, 2015

Abstract: Present study reports the (A) Clinical factors associated with DUB including 1) Incidence in different age groups. 2) Influence of parity in DUB. 3) Pattern of bleeding seen in DUB. 4) Influence of tubectomy, recent delivery and IUCD insertion on the incidence of DUB. 5) Association of DUB with medical diseases like diabetes and hypertension and (B) Identify the histopathology of endometrial curettage in cases of DUB.

The study is of prospective (observational) type. All patients of adolescent age group, reproductive and perimenopausal age group with history suggestive of dysfunctional uterine bleeding registered as inpatient and outpatient during the period August 1, 2010 to July 31, 2011 were studied. A total of 143 cases were selected. DUB forms about 10% of admissions in Government Medical College Hospital, Thrissur. Out of 156 cases, 143 cases were finally taken for the study. The commonest endometrial abnormality was proliferative endometrium. Anovulatory bleeding constituted 70% and ovulatory bleeding 30%. Uterine curettage is an important tool as depending on the functional status of the endometrium, treatment can be planned.

Key words: Abnormal uterine bleeding, Dysfunctional uterine bleeding, Ovulatory, anovulatory, Endometrial histology.

INTRODUCTION

Abnormal uterine bleeding affects 10-30 percent of reproductive aged women and upto 50 percent of perimenopausal women. Factors that impact the incidence most greatly are age and reproductive status.

Abnormal uterine bleeding can be due to the following:

1. Organic Causes
 - a. Diseases of the genital tract
 - b. Systemic diseases
2. Dysfunctional uterine bleeding.
 - a. Diseases of the Genital tract
 - i. Pregnancy-related conditions.
 - ii. IUCD - related bleeding
 - iii. Benign conditions-Endometriosis, Adenomyosis, Fibroids, Polyps
 - iv. Malignant lesions of the genital tract - Endometrial, Cervical, Vaginal.
 - b. Systemic Diseases-Coagulation disorders, Endocrinological abnormalities and liver disorders.

Dysfunctional uterine bleeding is one of the most frequently encountered problems in gynaecological practice.

It makes up about 10% cases in the outpatient clinic in the United States. It occurs most frequently at the extremes of menstrual life but it can develop at any intervening time.

Dysfunctional uterine bleeding is best defined as abnormal bleeding from the uterus in absence of organic disease of the genital tract, pregnancy or

general bleeding disorder. The term applies to any abnormal uterine bleeding, including disturbances of the menstrual cycle, regular and irregular uterine bleeding and alterations in the amount or duration of menstrual loss. But it most commonly implies excessive regular menstrual bleeding or essential menorrhagia (Van Eijkeren *et al.*, 1989). Diagnosis is one of exclusion.

It is generally accepted that a detailed history, careful general and pelvic examination, cervical smear, uterine curettage, hysteroscopy, or at least an endometrial biopsy are essential to exclude organic disease. There will nevertheless be a small proportion in whom some organic pathology is missed.

Thus DUB should always be regarded as a provisional diagnosis unless some endocrine or other dysfunction has been clearly demonstrated.

MATERIALS AND METHODS

The study is of prospective (observational) type. All patients of adolescent age group, reproductive and perimenopausal age group with history suggestive of DUB registered as in patient and outpatient during the period August 1, 2010 to July 31, 2011 were studied. A total of 143 cases were selected.

All cases where a pervaginal examination failed to reveal any recognizable pelvic pathology were included. All cases were subjected to cervical smear, ultrasound scan and selected after excluding organic pathology. Uterine size upto 6 weeks, small ovarian cysts and prolapsed uterus cases were not

*Corresponding Author:

Dr. Seena KB,
Assistant Professor,
Department of Obstetrics & Gynecology,
Government Medical College Hospital,
Thrissur, Kerala, India.



contraindications. Cases with obvious lesions of the cervix were excluded. Post-menopausal bleeding cases were also excluded.


The clinical detail of each patient was collected based on a preset proforma. Routine blood and urine examination and complete haemogram to rule out bleeding diathesis in the pubertal age group were carried out.

All selected cases were subjected to uterine curettage, 40% of the cases done during premenstrual and rest in bleeding phase. Curettings were sent for histopathological examination and reports were tabulated.

Appropriate treatment was advised to each case. Statistical analysis was done using the chi-squared test for goodness of fit and z-test for proportion.

RESULTS

156 cases with a clinical diagnosis of dysfunctional uterine bleeding were selected after subjecting them to ultrasound scan. 3 cases were due to IUCD use, 10 cases were excluded as they were found to have other pathologies, after curettage. Finally 143 cases were taken. Of these there were 5 cases of puberty menorrhagia, 10 cases with severe anaemia for whom hormonal treatment was given without subjecting them to curettage and 5 cases where there were no endometrial curettings.




1 ---<=20 years, 2----21-30 years,3 -----31-40 years,4-41-50 years,5---->=51 years

Age Pattern in the Study Group

The graph shows that the older age groups (41-50 yrs) are commonly affected, 65 cases (45.5%). Ages ranged from 12 yrs to 53 yrs. Cases of puberty menorrhagia constitute about 3.5% of the study group, 5 cases

PARITY & DUB



Parity ranged from nullipara to para. There were 10 nulliparous cases. Multiparous mostly para2 were commonly affected 60 cases, 41.96%.

Menorrhagia was the most common complaint in 84 cases (58.74%) followed by polymenorrhea 20 cases (13.99%) and metrorrhagia 17 cases (13.33%). Applying the chi-square test for goodness of fit the observed distribution is statistically significant. ($\chi^2 = 227.8$ $p < 0.001$)

Pattern of Bleeding In Each Age Group

MENORRHAGIA

AGE GROUP	NUMBER	PERCENTAGE
≤ 20 years	4	4.76%
21-30 years	10	11.9%
31-40 years	23	27.38%
41-50 years	40	47.6%
>51 years	7	8.33%


TOTAL = 84

Out of the total cases of menorrhagia, maximum (47.6%) was in the age group 41-50 years, followed by the age group 31-40 years (27.38%).

$\chi^2 = 70.51$ $P < .001$

Which is highly significant.

Out of the total cases of metrorrhagia, maximum number of cases was in the 31-40 age group. $\chi^2 = 6.8$ $P < .001$. Bleeding following period of amenorrhea was maximum in the 41-50 age group (71.43%). $\chi^2 = 12.29$ $p < 0.015$ Which is significant. There were 2 cases of polymenorrhagia and oligomenorrhagia both in the 41-50 age group. Out of the 20 cases of polymenorrhagia, maximum incidence was in the age group 31-40 years. $\chi^2 = 50.25$ $p < .001$ Which is significant. Maximum incidence of hypermenorrhagia was in 41-50 years. $\chi^2 = 9.5$ $p < .05$. There were 10 nullipara including 5 cases of puberty menorrhagia. Majority of the cases occurred in cases of last child birth ≥ 5 years which is natural as the age group most commonly affected is 41-50 years.




There were 80 cases that underwent puerperal sterilization and 21 cases that underwent interval sterilization.

Relation with IUCD

Menorrhagia 3
TOTAL = 3

In a total of 3 cases the abnormal bleeding was due to insertion of copper containing IUCDS. On removal of the device the bleeding pattern was normalized. All 3 of the cases with bleeding due to IUCD presented with menorrhagia and had Hb in the range 8-10. 2 of the cases belonged to the age group 21-30 years, and 1 in the 31- 40 year age group.

Anaemia in DUB



Majority of the cases, 114 had Hb in the range 8-10gm%. Severe degree of anemia Hb<5 gm%, was found in 2.8%. 4 cases of puberty menorrhagia had Hb 8-10 gm% and 1 in the group 5-7 gm%.


Histology of Endometrium in DUB Cases

Histopathology Of Endometrium	Number	Percentage
Secretory	36	29.27
Disordered Proliferative	21	17.07
Lytic	1	0.81
Proliferative	48	39.02
Complex Hyperplasia	5	4.06
Simple Hyperplasia	5	4.06
Irregular Ripening	1	0.81
Atypical Hyperplasia	6	4.87
Endometritis	0	0

TOTAL = 123

Proliferative endometrium formed the maximum number of cases (48) (39.02%) followed by secretory endometrium (36) (29.27%).

HISTOLOGY OF ENDOMETRIUM IN DUB CASES



Distribution of Endometrial Lesions In Various Age Groups

TYPE	<40 YEARS		> 40 YEARS		Z - VALUE
Proliferative Em.	18	32.14	30	44.78	1.44 (P>.05)
Secretory Em	29	51.79	7	10.45	5.04(P<.001)
Disordered Prolif. Em	6	10.71	15	22.39	1.42 (P>.05)
Lytic Em.	-	-	1	1.49	
Simple Hyperplasia	3	5.36	2	2.99	
Complex Hyperplasia	-	-	5	7.46	
Irregular Ripening	-	-	1	1.49	
Atypical Hyperplasia	-	-	6	8.96	
Endometritis	-	-	-	-	

TOTAL = 56 TOTAL = 67
 $\chi^2 = 32.7\%$ $P < .001$

The chi square statistics was found to be highly significant. This indicates differential distribution of endometrial lesions in the two age groups. Proliferative, disordered proliferative and hyperplastic endometrium were more in the above 40 age group, whereas secretory endometrium was more common in the below 40 year age group. The z test for proportion was applied and the above observation of secretory endometrium in below 40 was found significant.

Distribution of Normal Endometrium

Age Group	Number
< 40 YEARS	47
> 40 YEARS	37

TOTAL = 84

Normal endometrium was found in 47 cases in below 40 group and 37 cases in the above 40 group.

Relation between Abnormal Endometrium (Hyperplastic) and Age

Type	> 40 YEARS		< 40 YEARS	
	Number	Percentage	Number	Percentage
Disordered Prolif Em.	15	53.5	6	66.6
Simple Hyperplasia	2	71.4	3	33.33
Complex Hyperplasia	5	17.8	-	-
Atypical Hyperplasia	6	21.43	-	-
TOTAL = 28		TOTAL = 9		

Hyperplastic endometrium was most common in the above 40 age gr

Distribution of Ovulatory and Anovulatory Bleeding AGE GROUP


Type	> 40 Years		< 40 Years	
	Number	Percentage	Number	Percentage
Ovulatory	29	51.79	8	12.12
Anovulatory	27	48.21	58	87.88

Anovulatory bleeding was more common in above 40 age group. $X^2 = 42.38$ $P < 0.001$ highly significant.

Medical Diseases Associated With DUB

There were 8 cases of diabetes and 7 cases of hypertension in the study group.

TOTAL = 143



Curettage along with NSAIDS to control the bleeding was given in 76 cases. Curettage with hormonal treatment (combination OCPs or progesterone) was given for 34 cases. 10 cases presenting with heavy bleeding were treated with hormones, blood transfusion and hematinics. Symptomatic treatment was given for 5 cases of puberty menorrhagia with blood transfusion, hematinics and NSAIDS.

Hysterectomy In DUB

Hysterectomy, 3 as total and 15 as total abdominal with bilateral salpingo oophorectomy were

done. The cases included 6 cases of atypical hyperplasia and 12 cases who had recurrence of symptoms inspite of hormonal treatment and opted for hysterectomy.

DISCUSSION

Dysfunctional uterine bleeding is a common gynecological problem. Careful study of the endometrium is necessary especially in the older age group for, in a few, precursors of endometrial carcinoma will be found. Uterine curettage is useful since treatment can be planned depending on the histology and several unnecessary hysterectomies can be avoided.

REFERENCES

1. Wathen PI, Henderson, Witz CA Abnormal uterine bleeding Med clin NA 1995 Mar 79 (2) 329-44 (medline)
2. Falcone T, Desjardins C., Bourque J. Dysfunctional uterine bleeding in adolescents J reprod med 1994 oct 39 (10) 761-4 (medline)
3. John T Queenan Article on DUB Feb 2005 e-medicine page 1-17
4. Dallenbach-Hellweg Histopathology of endometrium 4th edition (1987) Berlin Springer-Verlag p-136.
5. Nedos S. DUB-Relation of endometrial histology to outcome, AmJO &G 109 (1971), 103-7.
6. Classification of endometrial hyperplasia (Ferenczy and Bergeson 1992) Fox text book of gynaec pathology chap 13-432.
7. Kurman Kaminski, Norris RJ. Endometrial hyperplasia-long term study of 170 patients- Cancer, 1985, 56,403-412.
8. Dysfunctional Uterine Bleeding An update. FOGSI publication, chapter9, 95-108.
9. MC Bride-Pre menopausal cystic hyper plasia and endometrial carcinoma. JO & G Brit. Emp. 66,288.
10. Ghosh and Sengupta KP-A study of the endometrium and cytohormonal pattern in DUB.J O&G India 1968, 18, 311.

Cite this article as:

Seena KB, A Study of Dysfunctional Uterine Bleeding: Clinical Factors and Endometrial Histology, International Journal of Bioassays, 2015, 4 (04), 000-000.

Source of support: Nil

Conflict of interest: None Declared