



Levels of serum zinc and manganese among post-menopausal patients with thyroid dysfunction

Sharique Ahmad^{1*}, Saeeda Wasim²

¹Department of Pathology, Era's Lucknow Medical College & Hospital, Lucknow, Uttar Pradesh, India.

²Department of Obstetrics & Gynecology, Integral Institute of Medical Sciences & Research, Lucknow, Uttar Pradesh, India.

Received for publication: December 28, 2015; Accepted: January 22, 2016

Abstract: The objective of this study is to shed more light on the role of trace metals and their mode of action in hyperthyroidism and hypothyroidism in Post-Menopausal Patients. The content of the trace elements Zinc and Manganese (Zn, and Mn) in the serum of patients was determined. Forty patients with hyperthyroidism and hypothyroidism were participated in this study. Serum zinc and manganese were determined using flame atomic absorption spectrophotometer technique. The study showed that serum zinc and manganese levels of hyperthyroidism patients were significantly lower ($p < 0.05$). While a significant increase in serum manganese level with decrease serum zinc levels in patients with hypothyroidism ($p < 0.05$). The present study confirmed a significant change in the levels of serum Zn, Mn in hyperthyroidism and hypothyroidism among Post-Menopausal patients and these changes may be related to pathophysiology of thyroid disease.

Key words: Hyperthyroidism; Hypothyroidism; Post-menopausal; Zinc; Manganese.

Introduction

The maintenance of optimal health requires an adequate supply of carbohydrates, proteins, lipid, and macronutrients, micronutrients and trace elements⁽¹⁾. Many trace elements play an essential role in number of biological processes through their action as activators or inhibitors of enzymatic reactions, by competing with other elements and proteins for binding sites, by influencing the permeability of cell membranes, or through other mechanisms. Trace elements are known to influence hormone at level of action, including hormone secretion and activity and binding to target tissue. Conversely, hormones influence trace metals metabolism at several levels of action, including excretion and transport of trace metals.⁽²⁾

Hence, trace elements assay in biological fluids can be used as diagnostic or prognostic aid in patients with different hormonal disturbances alongside with other biochemical parameters. Thyroid hormones regulate the rate of metabolic processes and consequent development of organism; deficiency of thyroid hormones causes many metabolic processes to slow down⁽³⁾.

Zinc is an important element for numerous biochemical processes as well as for cell proliferation. Zinc is extensively studied by bioinorganic chemistry and it is known that there are many metalloproteins with specific enzymatic activity contain zinc. Carbonic anhydrase, liver alcohol dehydrogenase and alkaline phosphatase are some examples of zinc enzyme⁽⁴⁾

Prasad *et al.*, (1990) reported that thyroid hormones modulate zinc transport activity in rate

renal and intestinal brush-border membrane.⁽⁵⁾ Zinc has been shown to have an antioxidant effect and stabilized cell membrane.⁽⁶⁾ Zinc is a metal that affects thyroid hormone function at several levels. For example, zinc deficiency inhibits TRH synthesis,⁽⁷⁾ and depresses plasma TSH, T4, T3 (triiodothyronine).⁽⁸⁾ It is necessary for extra thyroidal T4 to T3 conversion,⁽⁹⁾ and it plays a role in T3 binding to nuclear receptor as well as the binding of the receptor to DNA⁽¹⁰⁾.

Thyroid hormone receptors require zinc ion,⁽¹¹⁾ which facilitate folding into active shape⁽¹²⁾. There are indications that zinc is also important for normal thyroid homeostasis. Its roles are complex and may include effects on both the synthesis and mode of action of the hormones. Thyroid hormone binding transcription factors, which are essential for modulation of gene expression, contain zinc bound to cysteine residues.⁽¹³⁾ However, it is not known whether dietary zinc deficiency has a direct effect on this aspect of thyroid hormone metabolism. In cultured cells, very strong chelators of zinc are required to influence binding of transcription factors to DNA. In the thyroid gland itself, transcription factor 2, which interacts with the promoters for the thyroglobulin and thyroperoxidase genes, is a zinc-containing protein. The binding of transcription factor 2 is affected by redox state, but again it is not known whether this can be changed by dietary zinc intake.⁽¹⁴⁾

Zinc is very important in Thyrotropin-releasing hormone synthesis,⁽¹⁵⁾ is essential for thyroxine (T4)-to-T3 conversion,⁽¹⁶⁾ and is required for the

*Corresponding Author:

Dr. Sharique Ahmad,

Associate Professor,

Department of Pathology,

Era's Lucknow Medical College & Hospital,

Lucknow, Uttar Pradesh, India.

biological functioning of the thyroid hormone and related receptors.⁽¹⁷⁾

Manganese may affect thyroid hormone homeostasis and neurodevelopmental processes as a result of both direct dysregulation at the level of the thyroid gland and thyroid hormones, or indirectly via alterations in dopaminergic control of the thyroid gland and its hormones. Dopamine is a known modulator of both TSH and TSH subunit secretion.⁽¹⁸⁾ An additional effect of manganese on thyroid hormones homeostasis may be mediated through their metabolizing enzymes. Current data suggests that manganese can affect thyroid hormones directly by regulating the deiodinase enzymes.⁽¹⁹⁾

Materials and Methods

Patients:

Forty patients with hyperthyroidism and hypothyroidism were participated in this study. The patients diagnosed depending on the results of the following examinations: Clinical examinations, serum hormones (T3, T4) and TSH.

Samples collection and preparation:

A volume of 3 ml of venous blood sample was collected from both cases using sterile disposable syringe and aseptic standard vein puncture technique was applied, and emptied in a sterile plain container. And then sample was centrifuged at 4000 rpm for 10 minutes, then the serum was separated and transferred in a plain container and processed. Serum were separated and stored at -20°C until analysis.

Analysis of Trace Elements

Determination of Zinc and Manganese: Serum zinc and manganese were determined using flame atomic absorption spectrophotometer. Samples were diluted with deionized water. The analysis was performed against standard prepared in glycerol to approximate the viscosity characteristics of diluted samples. For determination of serum zinc, samples were diluted 1:5 with deionized water. The serum samples for manganese estimation were diluted 1:10.

Statistical Analysis:

The results of serum element concentrations were expressed as mean \pm standard deviation. The level of significance was determined by employing pooled t-test. Only when p-value less than 0.05, considered as statistically significant.

Results

Table 1 showed the results of serum trace elements expressed as (mean \pm standard deviation). Serum zinc and manganese levels of hyperthyroidism patients are significantly low. A significant decrease in serum zinc level was

demonstrated in patients with hypothyroidism while there is significant increase in serum manganese level.

Table 1: The serum concentration of trace elements in hypothyroidism, hyperthyroidism patients and reference values.

Serum Trace Elements	Hyperthyroidism	Hypothyroidism
Zinc mg/l (M \pm SD)	0.19 \pm 0.12	0.17 \pm 0.07
Manganese mg/l (M \pm SD)	0.15 \pm 0.11	1.72 \pm 1.36

Reference range of serum zinc level is (0.5 – 1.2, mean 85 mg/l) and manganese is (0.3 – 0.9, mean 0.6 mg/l).

Discussion

The present study showed significantly decrease in zinc and manganese content in serum of hyperthyroidism. This result is agreement with results of previous works performed on hyperthyroidism patients.⁽²⁰⁾

Zinc effects on thyroid hormones are complex and include both synthesis and mode of action.⁽²¹⁾ Thyroid transcription factors which are essential for modulation of gene expression contain zinc at cysteine residues.⁽¹²⁾

Manganese may affect thyroid hormone homeostasis and neurodevelopmental processes as a result of both direct dysregulation at the level of the thyroid gland and thyroid hormones, or indirectly via alterations in dopaminergic control of the thyroid gland and its hormones. Dopamine is a known modulator of both TSH and TSH subunit secretion.⁽¹⁷⁾

An additional effect of manganese on thyroid hormones homeostasis may be mediated through their metabolizing enzymes. Current data suggests that manganese can affect thyroid hormones directly by regulating the deiodinase enzymes.⁽¹⁸⁾ This finding disagree with observation reported by other workers that showed no significant difference in manganese concentration in serum of patients with hyperthyroidism.⁽²²⁾

The significant decrease in the level of zinc in hypothyroidism patients are observed in other different researches⁽²³⁾. One of possible explanation for this finding, that gastrointestinal absorption of zinc is severely impaired in hypothyroidism subjects. Low zinc level may reflect sequestration of zinc by liver or other tissues.⁽²⁴⁾ Another explanation is due to the significant influence of TSH in the variation of the concentration of human thyroid tissues.⁽²⁵⁾ Zinc has important roles in thyroid metabolism, It involves in T3 binding to its nuclear receptor, and participates in the formation and mechanism of action of TRH.⁽⁷⁾

A significant increase in manganese levels of hypothyroidism patients was observed. Manganese is a cofactor of many important enzymes especially, manganese superoxide dismutase, which is the principal antioxidant enzyme that neutralizes the toxic effects of reactive oxygen species. Manganese may directly or indirectly affect the thyroid function by injuring the thyroid gland or deregulating dopaminergic modulation of thyroid hormone synthesis.⁽¹⁷⁾

Conclusion

This study showed there is decrease in levels of zinc and manganese of hyperthyroidism patient while in hypothyroidism patient there is increase in serum manganese levels with decreasing in zinc levels. so, levels of trace elements like zinc and manganese should be evaluated in disturbed parameters of thyroid in co-relation with post-menopausal status of the patients.

References

1. Solomon's N Trace Elements in Clinical Nutrition: Parenteral Nutrition 2nd edition. Philadelphia, USA (1993) pp .150-183.
2. Stands bury J.B. and Kroc R.L. Human Development and Thyroid Gland: Relation to Endemic Cretinism, plenum press, New York (2000) p19.
3. Tapiero H. and Tew KD Trace elements in human physiology and pathology: zinc and metallothione ins. Biomedicine and pharmacotherapy (2003)57(9): pp399-411.
4. Bertini I The Coordination Chemistry of Metalloenzymes. The Role of Metals in reactions Involving water, oxygen, and related species. The Coordination Properties of the active site of zinc enzymes. In: Bertini I, Drago R S, Luchinat C, (eds), Kluwer Boston Inc. Canada. Published by Riedel Publishing Company.NA TO Advanced Study Institute, San Miniato-Italy (1982) pp1-3.
5. Prasad R, Kumar R, Singh KP Thyroid hormones modulate zinc transport activity of rate intestinal and renal brush-border membrane. Am J Physio (1999) pp1276: E774-782.
6. Powell Saul R. The antioxidant properties of Zinc. J Nutr (2000) pp 130:1447S-1454S.
7. Morley JE, Gordon J, Hershman JM. Zinc deficiency, chronic starvation, and hypothalamic-pituitary-thyroid function. Am J Clin Nutr. 1980 Aug;33(8): pp1767-70.
8. Pekary AE, Bhasin S, Smith V, Sugawara M, Swerdloff RS, Hershman JM. Thyroid hormone modulation of thyrotrophin-releasing hormone (TRH) and TRH-Gly levels in the male rat reproductive system. J Endocrinol. 1987 Aug;114(2): pp271-7.
9. Pekary AE, Lukaski HC, Mena I, Hershman JM. Processing of TRH precursor peptides in rat brain and pituitary is zinc dependent. Peptides. 1991 Sep-Oct;12(5):pp1025-32.
10. Anselmet A, Bismuth J, Torresani J. Triiodothyronine nuclear receptor. Role of histones and DNA in hormone binding. Biochim Biophys Acta. 1983 Apr 15; pp 739(3):291-300.
11. Sustrova M, Strbak V. Thyroid function and plasma immunoglobulins in subjects with Down's syndrome (DS) during ontogenesis and zinc therapy. J Endocrinol Invest. Jun (1994) 17(6): pp 385-90.
12. Bucci I, Napolitano G, Giuliani C, *et al.*, Zinc sulphate supplementation improves thyroid function in hypozincemics Down children. Biol Trace Elem Res.; (1999)67: pp 257-268.
13. Civitareale D, Saiardi A, Falasca P. Purification and characterization of thyroid transcription factor 2. Biochem 1994; pp 304: 981-5
14. Kralik A, Eder K, Kirchgessner M. Influence of zinc and selenium deficiency on parameters relating to thyroid hormone metabolism. Horm Metab Res 1996; pp 28: 223-6.
15. Pekary AE, Lukaski HC, Mena I, Hershman JM. Processing of TRH precursor peptides in rat brain and pituitary is zinc dependent. Peptides 1991;12(5): pp 1025-32.
16. Chen MD, Lin PY, Lin WH. Zinc supplementation on serum levels and hepatic conversion of thyroid hormones in obese (ob/ob) mice. Biol Trace Elem Res 1998;61(1): pp 89-96.
17. Freake HC, Govoni KE, Guda K, Huang C, Zinn SA. Actions and interactions of thyroid hormone and zinc status in growing rats. J Nutr 2001;131(4): pp 1135-41.
18. Konig S, MouraNeto V. Thyroid hormone actions on neural cells. Cell Mol Neurobiol. 2002; pp 22:517-44.
19. Aihara K, Nishi Y, Hatano S, Kihara M, Yoshimitsu K, Tekeichi N, *et al.* Zinc, copper, manganese, and selenium metabolism in

- thyroid disease. *Am J Clin Nutr.* 1984; pp 40:26–35.
20. Vitoux, D., Arnaud, J., and Chappuis, P. " Are copper, zinc and selenium in erythrocytes valuable biological indexes in nutrition and pathology?" *J Trace Elem Med Biol.* 1999; pp 13,113-128.
 21. Arthur JR, Beckett GJ Thyroid function. *British Medical Bulletin* (1999) 55: pp 658-668.
 22. Ali, E.A; Tahssen, Y.H; Saleh, B.O.M Study of some Trace elements in hyperthyroidism patients (2007) 6 NO 2, pp 71-79
 23. Buchinger W, Leopold B, Lind P, Langsteger W, Klima G, Koltringer P, et al Changes in zinc level in serum, whole blood and erythrocytes in disorders of thyroid function. *Wien Klin Wochenschr* (1988)100 (18): pp 619-21.
 24. Yoshida k, kiso Y, Watanabe T, Kaise, K, Kaise N, and ItagaKi M Erythrocyte zinc in hyperthyroidism: reflection of integrated thyroid hormone levels over the previous few months. *Metabolism* (1990) 39(2): pp 182-186.
 25. Bellisola G, Bratter P, Cinque G, Francia G, Galassini S, Gawlik D, et al., The TSH-dependent variation of the essential elements iodine, zinc and selenium within human thyroid tissues *Journal of Trace Elements in Medicine and Biology* (1998) pp 12(3): 177-82.

Cite this article as:

Sharique Ahmad and Saeeda Wasim. Levels of Serum Zinc and Manganese among Post-Menopausal Patients with Thyroid Dysfunction. *International Journal of Bioassays* 5.2 (2016): 4821-4824.

Source of support: Nil

Conflict of interest: None Declared